

DETECTION OF *PSEUDOMONAS AVELLANAE* FROM HAZELNUT TWIGS BY TAQMAN REAL-TIME PCR

F. Gervasi and M. Scortichini

C.R.A.-Unità di Ricerca per la Frutticoltura, Via Torrino 3, 81100 Caserta, Italy

SUMMARY

A TaqMan real-time PCR assay was developed to detect *Pseudomonas avellanae*, the causal agent of bacterial canker and decline of hazelnut (*Corylus avellana* L.). The real-time primers and probe were designed within the 16S rRNA gene of the pathogen after a comparison with other related species and pathovars. The assay detected DNA from a panel of *P. avellanae* strains obtained from different geographic areas. The other bacterial pathogens inciting diseases to hazelnut, *P. syringae* pv. *coryli*, *P. s.* pv. *syringae* and *Xanthomonas arboricola* pv. *corylina*, did not react in the assay. Only *P. s.* pv. *actinidiae*, isolated from kiwifruit and yellow kiwifruit in Italy, cross-reacted. The assay detected the pathogen when inoculated in low numbers (80 cells) in a leaf scar up until nine days after inoculation. The addition of bovine serum albumin was essential to prevent inhibition by compounds released from the hazelnut tissue. This assay could be useful for monitoring the presence of *P. avellanae* either in orchards or nurseries. Screening for germplasm resistance could also benefit from this technique.

Keywords: hazelnut decline, real-time PCR, genomospecies, *Pseudomonas*, *Xanthomonas*.

INTRODUCTION

Pseudomonas avellanae (Psallidas) Janse *et al.*, the cause of bacterial canker and decline of hazelnut (*Corylus avellana* L.), provokes serious economic losses in central Italy (Viterbo province) and northern Greece (Scortichini, 2002). Identification and characterization of the agent is currently possible by using classical (biochemical tests, fatty acid analysis) and molecular (repetitive-PCR) techniques (Janse *et al.*, 1996; Scortichini *et al.*, 2002). For detection, two techniques were developed. Scortichini and Marchesi (2001) developed a

PCR using the 16S rRNA gene sequence. An amplicon of 762 bp was specifically produced only by the *P. avellanae* strains isolated in Italy and Greece and the sensitivity with pure cultures of the pathogen was up to 6.5×10^2 CFU ml⁻¹. Loreti and Gallelli (2002) designed specific primers directed to the *hrpW* gene of the bacterium and a PCR assay. An amplicon of 350 bp specifically detected all *P. avellanae* strains from Italy and Greece and the sensitivity with pure cultures reached 2.0×10^3 CFU ml⁻¹. Both techniques could detect the bacterium in infected plant material. It proved critical, however, for the effectiveness of the 16S rRNA-based protocol to add specific compounds in order to limit the negative effect that hazelnut polyphenolics exert on PCR amplification (Scortichini and Marchesi, 2001).

Conventional PCR assays can pose some limitations. They are not useful for quantitative analysis, need post-amplification analysis such as gel electrophoresis and UV visualization of products, and they are exposed to the risk of contamination. Real-time PCR, in contrast, can amplify and simultaneously quantify a target DNA without the need of post-amplification manipulation. It can be used in a high-throughput manner and it is suitable for large-scale monitoring and screening procedures. It has been successfully applied to detect pathogenic bacteria from plant tissue (reviewed by Palacio-Bielsa *et al.*, 2009).

In this study we report on the development of a TaqMan real-time PCR assay to detect *P. avellanae* from hazelnut twigs.

MATERIALS AND METHODS

Bacterial strains. The strains of *P. avellanae* and other bacteria used in this study are listed in Table 1. Pseudomonads and *Pantoea agglomerans* were grown on nutrient agar with 5% of sucrose added (NSA) at 25-27°C. Xanthomonads were cultured on glucose-yeast-extract-calcium-carbonate agar (GYCA).

DNA preparation. Bacterial DNA preparations were obtained starting from pure cultures grown for 48 h on the appropriate culture medium. A loopful of pure cul-

Table 1. List of bacterial species and pathovars used in this study and response of the TaqMan real-time assay.

Species/pathovar	Strain designation	Origin, year of isolation, host	Real-time PCR
<i>Pseudomonas avellanae</i>	BPIC 631 ^T	Greece, 1976, <i>Corylus avellana</i>	+
	CRA-PAV 013	Italy, 1992, <i>C. avellana</i>	+
	CRA-ISF 2059	Italy, 1994, <i>C. avellana</i>	+
	CRA-PAV 1265	Italy, 2003, <i>C. avellana</i>	+
	CRA-FRU NOC2	Italy, 2006, <i>C. avellana</i>	+
	CRA-FRU NOC3	Italy, 2006, <i>C. avellana</i>	+
	CRA-FRU VNOC3	Italy, 2007, <i>C. avellana</i>	+
	CRA-FRU VTGR9	Italy, 2007, <i>C. avellana</i>	+
<i>Pseudomonas syringae</i> pv. <i>coryli</i>	NCPPB 4273 ^T	Italy, 1995, <i>Corylus avellana</i>	-
	PD 3453	Italy, 1995, <i>C. avellana</i>	-
	CRA-PAV 598	Italy, 1995, <i>C. avellana</i>	-
	DPP 48	Italy, 1999, <i>C. avellana</i>	-
	DPP 51	Italy, 1999, <i>C. avellana</i>	-
<i>Pseudomonas syringae</i> pv. <i>morsprunorum</i>	NCPPB 2427	Switzerland, 1971, <i>Prunus armeniaca</i>	-
	NCPPB 2787	Greece, 1975, <i>Prunus avium</i>	-
<i>Pseudomonas syringae</i> pv. <i>syringae</i>	CRA-PAV 14a	Italy, 1998, <i>Corylus avellana</i>	-
	CRA-PAV P11	Italy, 1998, <i>C. avellana</i>	-
	CRA-PAV 1365	Italy, 2006, <i>C. avellana</i>	-
	CRA-PAV 1386	Italy, 2007, <i>C. avellana</i>	-
	CRA-PAV 1394	Italy, 2007, <i>C. avellana</i>	-
	CRA-PAV 1431	Italy, 2007, <i>C. avellana</i>	-
<i>Pseudomonas syringae</i> pv. <i>tomato</i>	CRA-FRU 228	Brasil, 1996, <i>Lycopersicon esculentum</i>	-
	CRA-FRU 617	Italy, 1996, <i>L. esculentum</i>	-
<i>Pseudomonas syringae</i> pv. <i>actinidiae</i>	NCPPB 3739 ^T	Japan, 1984, <i>Actinidia deliciosa</i>	-
	CRA-FRU Act 2	Italy, 2001, <i>A. deliciosa</i>	+
	CRA-FRU Act 3	Italy, 2001, <i>A. deliciosa</i>	+
	CRA-FRU Act 4	Italy, 2001, <i>A. deliciosa</i>	+
	CRA-FRU Act chi 1	Italy, 2008, <i>A. chinensis</i>	+
	CRA-FRU Act chi 2	Italy, 2008, <i>A. chinensis</i>	+
	CRA-FRU Act chi 3	Italy, 2008, <i>A. chinensis</i>	+
	CRA-FRU Act chi 4	Italy, 2008, <i>A. chinensis</i>	+
<i>Pseudomonas viridiflava</i>	NCPPB 1252	United Kingdom, 1960, <i>Pyrus communis</i>	-
<i>Pseudomonas corrugata</i>	NCPPB 2450	United Kingdom, 1972, <i>Lycopersicon esculentum</i>	-
	CRA-FRU 21	Italy, 1996, <i>L. esculentum</i>	-
<i>Pseudomonas marginalis</i> pv. <i>marginalis</i>	PD 1630	U.S.A., 1990, <i>Dieffenbachia</i> sp.	-
<i>Pseudomonas savastanoi</i> pv. <i>fraxini</i>	PD 117	The Netherlands, 1978, <i>Fraxinus excelsior</i>	-
<i>Xanthomonas arboricola</i> pv. <i>corylina</i>	NCPPB 935 ^T	India, 1959, <i>Corylus maxima</i>	-
	NCPPB 984	U.S.A., 1941, <i>C. maxima</i>	-
	NCPPB 2896	United Kingdom, 1976, <i>C. avellana</i>	-
	CRA-FRU 235	Italy, 1997, <i>C. avellana</i>	-
	CRA-FRU 238	Italy, 1999, <i>C. avellana</i>	-
	CRA-FRU 504	Italy, 2002, <i>C. avellana</i>	-
	CRA-FRU 505	Italy, 2005, <i>C. avellana</i>	-
CRA-FRU 3657	Italy, 2007, <i>C. avellana</i>	-	
<i>Pantoea agglomerans</i>	CRA-FRU 14aq	Italy, 2002, <i>Corylus avellana</i>	-

BPIC: Culture Collection of the Benaki Phytopathology Institute, Kiphissia-Athens, Greece

CRA-ISF: Culture Collection of Centro di Ricerca per la Frutticoltura, Roma, Italy

CRA-PAV: Culture Collection of Centro di Ricerca per la Patologia Vegetale, Roma Italy

DPP: Culture Collection of Department of Plant Protection, University of Sassari, Italy

NCPPB: National Collection of Plant Pathogenic Bacteria, York, United Kingdom

PD: Culture Collection of the Plant Protection Service, Wageningen, The Netherlands

ture was suspended in sterile bidistilled water and centrifuged at 13,000 *g* for 2 min. The pellet was resuspended in 100 μ l of 50 mM NaOH, and the suspension heated at 95°C for 15 min, and centrifuged 13,000 *g* for 2 min. The supernatant was collected, an equal volume of chloroform/isoamyl alcohol (24:1) added, mixed thoroughly, and centrifuged at 10,000 *g* for 5 min at room temperature. The aqueous phase was transferred to a new tube, 5 M NaCl added to a final concentration of 2 M NaCl, mixed with two volumes of absolute ethanol, and kept 2 h at -20°C. Then this phase was centrifuged at 14,000 *g* for 10 min at 4°C; the supernatant was decanted, the pellet washed with 70% (v/v) ethanol, air dried and dissolved in 40 μ l of double distilled water. The quantity and quality of the DNA were determined by spectrophotometer (GeneQuant, Pharmacia, USA).

Primer and probe design. The 16S gene sequences of *P. avellanae* and related pseudomonads from the NCBI databank were aligned by the BioEdit sequence alignment editor (Ibis Bioscience, USA). Within these alignments, specific regions, unique to each target organism, were identified. A BLAST search was performed in order to find the best consensus sequence and to ensure that no non-target organism of which a sequence is present in the database had a homologous sequence. Annealing sites for primers and the specific TaqMan probe were manually designed within the region exhibiting the best consensus sequence. Two primers and a probe were designed based on the regions of the 16S rRNA gene of *P. avellanae* ISF 2059 which exhibited suitable variability within the region when compared with the related pseudomonads (Table 2). The distance of the primers was chosen to cover a length of 82 bp to ensure efficient PCR replication. The forward primer, Pavel-F, was a 20-mer covering positions 422-441. The reverse primer, Pavel-R, was a 19-mer covering positions 486-504. The probe, Pavel-MGB, is a 17-mer TaqMan-MGB probe covering the positions 454-470, a region between that of the primers.

Primers and probe were synthesized by Applied Biosystems (USA). The minor groove binder (MGB)-probe was covalently labelled by the manufacturer at the 5' end with the reporter dye FAM and, at the 3' end, with the non-fluorescent quencher NFQ1. The MGB

probe increases the affinity of the TaqMan probe for the target sequence allowing shorter probes to be used, often increasing the sequence specificity of the detection system, and improving the quenching of the reporter signal due to the shorter distance between reporter and quencher. Moreover, the MGB probe allows the design of a probe with a low G/C percentage; this stabilizes the A/T rich duplexes more than the G/C ones increasing the T_m of A/T rich probes (Kutyavin *et al.*, 2000).

TaqMan real-time PCR assay conditions. The reactions were performed in 96-well plates in a Bio-Rad iCycler thermal cycler iQ5 (Bio-Rad, USA). Amplification reaction mixture (25 μ l) consisted of 2 μ l of DNA template, 1 \times PCR Buffer II (Applied Biosystems, USA), 4 mM MgCl₂, 200 μ M dNTPs (each), 300 nM primers (each), 200 nM TaqMan MGB probe, 1.25 U AmpliTaq Gold DNA polymerase (Applied Biosystems, USA). For detection in crude plant tissue extracts, 400 ng/ μ l ultra-pure bovine serum albumin (BSA) (Applied Biosystems, USA) was added to reduce the negative effect of polyphenols on PCR amplification (Kreader, 1996; Weller *et al.*, 2000; Kageyama *et al.*, 2003). The template was 10 ng/PCR of DNA extracted from pure culture for the assay specificity or 2 μ l from crude plant extract for spiking to test the assay *in vivo*. Cycling conditions were as follows: initial activation at 95°C for 10 min, followed by 45 two-step cycles comprising 10 sec at 95°C and 45 sec at 60°C.

Performance and specificity of the assay. Performance was analysed by running five ten-fold serial dilutions of purified *P. avellanae* CRA-PAV 1265 DNA in sterile double-distilled water. Two separate runs of two separate *P. avellanae* DNA dilution series were analysed. The mean slope of the linear regression line between logarithmic values of *P. avellanae* DNA concentration and cycle threshold number (Ct) values was used to calculate the amplification efficiency, $E=(10^{[-1/\text{slope}]})-1$, where 1 corresponds to 100% amplification efficiency (Pfaffl, 2001; Rasmussen, 2001). The correlation coefficient (R^2) was also determined. The specificity of the primers targeting *P. avellanae* was assessed by checking the DNAs of related bacterial species and pathovars listed in Table 1, and prepared as described above for *P. avellanae*. Each DNA was tested in dupli-

Table 2. Nucleotide sequences of TaqMan real-time PCR primers and of the minor groove binder (MGB) probe developed in this study.

Primer/Probe	Orientation	Sequence (5'-3')	Length (nt)	T_m (°C)	GC content (%)
<i>Pavel-F</i>	Forward	GCACTTTAAGTTGGGAGGAA	20	64.5	45
<i>Pavel-R</i>	Reverse	CAGAGTTAGCCGGTGCTTA	19	65.8	52.6
<i>Pavel-MGB</i>	Probe	AATACGTATCTGTTTTG-MGB	17	52.1	29.4

cate, and *P. avellanae* CRA-PAV 1265 was used as positive control.

Sensitivity of the assay for infected hazelnut twigs.

Samples of artificially inoculated hazelnut material were used. Twigs of field-cultivated hazelnut cultivar Tonda Gentile Romana were inoculated at the beginning of autumn by following the method of Scortichini and Lazari (1996) and Scortichini *et al.* (2002b). Ten μl of bacterial suspensions of *P. avellanae* 1265 serially diluted ten-fold in sterile physiological saline (SPS; 0.85% NaCl in distilled water) from 8×10^7 CFU ml^{-1} to 8×10^2 CFU ml^{-1} photometrically determined, were placed, at the beginning of October, on the leaf scar of a one-year old twig, using 30 twigs. Each dose was replicated five times. Control twigs were inoculated with SPS only. After one, two, seven and nine days post-inoculation, the twigs were removed from the tree and two slices (20-40 mg total) from the leaf scar were excised using a sterile scalpel, collected in a 2.0 ml Eppendorf tube, 320 μl of SPS added, and crushed by a sterile micro pestle. After an incubation of 25 min at room temperature, 280 μl of the suspension was collected into a new tube and centrifuged at 13,000 g for 10 min. The supernatant was discarded, and the pellet resuspended in 20 μl of SPS. Two μl of this crude suspension were directly used as template in the PCR. Each crude suspension was run in three replicates and the mean Ct values calculated.

RESULTS

Performance and specificity of the assay. Parameters were determined by amplifying ten-fold dilutions of *P. avellanae* DNA (from 100 ng to 10 pg per reaction). Each dilution was amplified in three replicates and the




Fig. 1. Linear regression generated by ten-fold dilution of bacterial DNA (picograms) and the threshold cycle values as detected by the TaqMan real-time PCR assay. The amplification efficiency is 99.7 and the regression R^2 is 0.99.

mean Ct used for a linear regression analysis. A slope of -3.33 (efficiency 99.7) and a regression R^2 value of 0.99 indicated a good fitness for a real-time PCR assay (Fig. 1). A panel of *P. avellanae* strains and other bacterial species were tested to check the specificity. Results are summarized in Table 1. The assay detected all *P. avellanae* strains from various geographic locations used in this study (Fig. 2). The amplification plot showed that all isolates crossed the threshold-line between cycles 14 and 15, when the same amount of bacterial DNA (from 100 ng to 10 pg per reaction per strain) was tested. The other hazelnut pathogens (i.e. *P. syringae* pv. *coryli*, *P. s.* pv. *syringae* and *X. a.* pv. *corylina*), other phytopathogenic pseudomonad strains, and *Pantoea agglomerans* did not react (Fig. 3). Remarkably, *P. s.* pv. *actinidiae* strains originating from Italy showed a strong signal with our probe. *P. s.* pv. *actinidiae* NCPPB 3739, the type-strain of the pathovar, isolated in Japan, however, did not show amplification. No signal was detected for the negative controls included in the test.

Sensitivity of the assay with samples of infected hazelnut twigs. Detection sensitivity was determined using three replicates of extracts of artificially inoculated hazelnut twigs. The assay was able to detect *P. avellanae* CRA-PAV 1265 artificially inoculated in low concentration (i.e. 8×10^3 CFU/ml) on/in leaf scars up to nine days after inoculation. The addition of BSA proved essential to enable DNA amplification since no amplification was recorded when it was not added. No amplification signal was detected for the negative controls.




Fig. 2. Sensitivity of the TaqMan real-time PCR assay for the detection of *Pseudomonas avellanae*. Amplification plot from the Bio-Rad iCycler thermal cycler iQ5 detection system showing cycle threshold number (Ct) versus normalized fluorescent values of a dilution series of DNA from a panel of *P. avellanae* strains. Different panel of curves represent *P. avellanae* strains from Italy and the type-strain from Greece tested with the assay at decreasing doses. The reaction did not produce any detectable amplicon for the negative control samples.




Fig. 3. Specificity of the TaqMan real-time PCR assay for the detection of *Pseudomonas avellanae*, for a number of other bacterial hazelnut pathogens, other phytopathogenic pseudomonads and *Pantoea agglomerans*. Curves showing fluorescence increasing at cycle threshold number (Ct) between 14 and 15 correspond to DNA preparations from pure cultures of *P. avellanae* strains from Italy and the type-strain from Greece and an Italian strain of *P. s. pv. actinidiae*. The reaction did not produce any detectable amplicon (i.e. no increasing curves present at the bottom of the plot) for all the other species and pathovars reported in Table 1, including the type strain of *P. s. pv. actinidiae* from Japan.

DISCUSSION

The assay developed has potential for rapid detection of *P. avellanae*. Primers and probe targeted to the 16S rRNA gene of the bacterium enabled detection of a panel of *P. avellanae* strains from Greece and Italy. None of the strains we tested of other bacterial pathogens inciting diseases on hazelnut, namely *P. s. pv. coryli*, *P. s. pv. syringae* and *X. a. pv. arboricola*, cross-reacted in the assay. Some Italian strains of *P. s. pv. actinidiae*, the causal agent of bacterial canker of kiwifruit (*Actinidia deliciosa* Liang et Ferguson) and yellow kiwifruit (*A. chinensis* Planchon) strains isolated in Italy (Scortichini, 1994; Ferrante and Scortichini, 2009), however, strongly cross-reacted with our probe. The *P. s. pv. actinidiae* type-strain, isolated in Japan (Takikawa et al., 1989), did not cross-react. These findings once more confirm the genetic relatedness of this pathovar with *P. avellanae* (Scortichini et al., 2002b) and point out that genetic variability exists within this pathovar. Similarly, Weller et al. (2007) found that their TaqMan real-time PCR assay, exploiting the *pep* (prolylendopeptidase) gene of *Xanthomonas arboricola*, detected not only *X. a. pv. fragariae* but also cross-reacted with some strains of *X. a. pv. corylina* and *pv. pruni*.

For reliable use of the protocol, addition of BSA proved critical since, as previously noted (Scortichini and Marchesi, 2001), PCR-inhibiting compounds are present in macerated hazelnut twigs. For preliminary adjustment of the assay, we monitored the pathogen up

to nine days from inoculation. Additional study is necessary to further investigate the sensitivity and specificity of the real-time PCR assay in for use in epidemiological studies with epiphytic and/or latent populations of the pathogen.

REFERENCES

- Ferrante P., Scortichini M., 2009. Identification of *Pseudomonas syringae* pv. *actinidiae* as causal agent of bacterial canker of yellow kiwifruit (*Actinidia chinensis* Planchon) in central Italy. *Journal of Phytopathology* **157** (in press).
- Kageyama K., Komatsu T., Suga H., 2003. Refined PCR protocol for detection of plant pathogens in soil. *Journal of General Plant Pathology* **69**: 153-160.
- Kreader C.A., 1996. Relief of amplification inhibition in PCR with bovine serum albumin or T4 gene 32 protein. *Applied and Environmental Microbiology* **62**: 1102-1106.
- Kutyavin I.V., Afonina I.A., Mills A., Gorn V.V., Lukhtanov E.A., Belousov E.S., Singer M.J., Walburger D.K., Lokhov S.G., Gall A.A., Dempcy R., Reed M.W., Meyer R.B., Hedgpeth J., 2000. 3' minor groove binder-DNA probes increase sequence specificity at PCR extension temperatures. *Nucleic Acids Research* **28**: 655-661.
- Janse J.D., Rossi M.P., Angelucci L., Scortichini M., Derks J.H.J., Akkermans A.D.L., De Vrijer R., Psallidas P., 1996. Reclassification of *Pseudomonas syringae* pv. *avellanae* as *Pseudomonas avellanae* (spec. nov.), the bacterium causing canker of hazelnut (*Corylus avellana* L.). *Systematic and Applied Microbiology* **19**: 589-595.
- Loreti S., Gallelli A., 2002. Rapid and specific detection of virulent *Pseudomonas avellanae* strains by PCR amplification. *European Journal of Plant Pathology* **108**: 237-244.
- Palacio-Bielsa A., Cambra M.A., López M.M., 2009. PCR detection and identification of plant-pathogenic bacteria: updated review of protocols (1989-2007). *Journal of Plant Pathology* **91**: 249-297.
- Pfaffl M.W., 2001. A new mathematical model for relative quantification in real-time RT PCR. *Nucleic Acids Research* **29**: 2002-2007.
- Rasmussen R., 2001. Quantification in the lightcycler. In: Meuer S., Wittwer C., Nakagawara K. (eds). Rapid cycle real-time PCR. Methods and Applications, pp. 21-34. Springer, Heidelberg, Germany.
- Scortichini M., 1994. Occurrence of *Pseudomonas syringae* pv. *actinidiae* in Italy. *Plant Pathology* **43**: 1035-1038.
- Scortichini M., Lazzari M., 1996. Systemic migration of *Pseudomonas syringae* pv. *avellanae* in twigs and young shoots of hazelnut and symptom development. *Journal of Phytopathology* **144**: 215-219.
- Scortichini M., Marchesi U., 2001. Sensitive and specific detection of *Pseudomonas avellanae* using primers based on 16S rRNA gene sequences. *Journal of Phytopathology* **149**: 527-532.
- Scortichini M., 2002. Bacterial canker and decline of European hazelnut. *Plant Disease* **86**: 704-709.
- Scortichini M., Marchesi U., Rossi M.P., Di Prospero P., 2002a. Bacteria associated with hazelnut (*Corylus avellana* L.) decline are of two groups: *Pseudomonas avellanae* and

- strains resembling *P. syringae* pv. *syringae*. *Applied and Environmental Microbiology* **68**: 476-484.
- Scortichini M., Marchesi U., Di Prospero P., 2002b. Genetic relatedness among *Pseudomonas avellanae*, *P. syringae* pv. *theae* and *P. s.* pv. *actinidiae*, and their identification. *European Journal of Plant Pathology* **108**: 269-278.
- Takikawa Y., Serizawa S., Ichikawa S., Tsuyumu S., Goto M., 1989. *Pseudomonas syringae* pv. *actinidiae* pv. nov., the causal bacterium of canker of kiwifruit in Japan. *Annals of the Phytopathological Society of Japan* **54**: 224-228.
- Weller S.A., Beresford-Jones N.J., Hall J., Thwaites R., Parkinson N., Elphinstone J.G., 2007. Detection of *Xanthomonas fragariae* and presumptive detection of *Xanthomonas arboricola* pv. *fragariae*, from strawberry leaves, by real-time PCR. *Journal of Microbiological Methods* **70**: 379-383.

Received February 6, 2009

Accepted April 27, 2009